

M. B. Black · K. M. Halanych · P. A. Y. Maas
 W. R. Hoeh · J. Hashimoto · D. Desbruyères
 R. A. Lutz · R. C. Vrijenhoek

Molecular systematics of vestimentiferan tubeworms from hydrothermal vents and cold-water seeps

Received: 11 July 1997 / Accepted: 21 August 1997

Abstract Vestimentiferan tubeworms inhabit sulfide-rich environments associated with deep-sea hydrothermal vents and cold-water seeps at continental margins. Twelve species have been described, and several more await formal descriptions. As a group, these worms are best known for their lack of a digestive system in adults and their dependence on endosymbiotic bacteria that supply nutrients derived from chemoautotrophism. The taxonomic status of Vestimentifera has been debated since their discovery. Furthermore, relationships within the Vestimentifera have been difficult to determine by morphological criteria. Several species display considerable phenotypic plasticity, further confounding efforts to establish evolutionary relationships. We used a fragment of the mitochondrial cytochrome c oxidase subunit I gene to examine evolutionary relationships among vent-endemic species (*Riftia pachyptila*, *Oasisia alviniae*, *Ridgeia piscesae*, *Tevnia jerichonana*) and seep-associated species (*Escarpia laminata*, *E. spicata*, *Lamellibrachia barhami*, *L. columna*, and an undescribed species) of these worms. The molecular data placed these vestimentiferan taxa within the phylum Pogonophora. The

pogonophoran clade (including vestimentiferans) was then linked to the Annelida. Examination of sequence divergence suggests that extant vestimentiferans constitute a recent evolutionary radiation that diversified as a paraphyletic assemblage of seep-associated taxa (including the genera *Lamellibrachia* and *Escarpia*) and then gave rise to a clade of vent-endemic taxa (genera *Riftia*, *Oasisia*, *Ridgeia* and *Tevnia*).

Introduction

Vestimentiferan tubeworms are among the dominant fauna associated with hydrothermal vents in the Pacific Ocean. They also occur in sulfide-rich cold seeps along continental margins. As adults they have no mouth, gut, or anus, and depend on sulfur-oxidizing bacterial endosymbionts for nutriment (Cavanaugh et al. 1981; Felbeck 1981). Webb (1969) erected the order Vestimentifera to encompass these worms, which he considered as members of the class Pogonophora (Phylum: Annelida). Descriptions of new vestimentiferan species quickly followed the discovery of hydrothermal vents in the eastern Pacific. In revising the taxonomy of these unusual worms, Jones (1985) recommended the Vestimentifera be elevated to phylum level and considered a sister taxon to the phylum Pogonophora (cf. Mañé-Garzón and Montero 1985; Jones 1987).

Southward (1988, 1991, 1993) does not recognize Jones' (1985) new phylum. She treats vestimentiferans as a discrete class (the Obturata) of the phylum Pogonophora. Earlier, Jones (1981) had also recognized vestimentiferans as members of the Pogonophora. He placed vestimentiferans in a new subphylum, the Obturata, and relegated all other pogonophorans to the subphylum Perviatea. Based on morphology, vestimentiferans and perviate pogonophorans have been linked to the Annelida (Nørrevang 1970, 1974; Van der Land and Nørrevang 1975; Bartolomaeus 1995). Some authors (Webb 1964; Ivanov 1988; Southward 1988; Jones and Gardiner 1989) base this relationship partly

Communicated by N.H. Marcus, Tallahassee

M.B. Black¹ · K.M. Halanych · P.A.Y. Maas ·
 W.R. Hoeh² · R.A. Lutz · R.C. Vrijenhoek (✉)
 Center for Theoretical and Applied Genetics
 and Institute of Marine and Coastal Sciences,
 Rutgers University, New Brunswick,
 New Jersey 089903-0231, USA

J. Hashimoto
 Japan Marine Science and Technology Center (JAMSTEC),
 2-15 Natsushima-Cho, Yokosuka 237, Japan

D. Desbruyères
 URM No. 7, Département Environnement Profond,
 IFREMER, BP 70, F-29280 Plouzané Cedex, France

Present addresses:

¹ Maurice Lamontagne Institute, PO Box 1000,
 Mont-Joli, PQ.G5H 3Z4, Canada

² Department of Zoology, Miami University, Oxford,
 Ohio 45056, USA

on a free-feeding trochophore-like larval stage, but this primitive characteristic is shared by many metazoan taxa. The association with annelids remains inconclusive (Rouse and Fauchald 1995). Ivanov (1988, 1994) treats the Pogonophora (incorporating vestimentiferans) as distinct from the phylum Annelida. Crucial questions remain, however, regarding development of the coelom, the nervous and circulatory systems, and segmentation in pogonophorans and vestimentiferans (Southward 1993).

Molecular systematics offers an alternative approach to inferring evolutionary relationships among taxa that have proven difficult to resolve with morphological and developmental studies. Recent studies involving mitochondrial cytochrome oxidase I and nuclear ribosomal sequences support a close affiliation between vestimentiferans and perviate pogonophorans (Winnepenninckx et al. 1995). However, the exact placement of a vestimentiferan and pogonophoran clade among protostome metazoans is questionable. Although some studies ally these organisms with annelids (Field et al. 1988; Suzuki et al. 1988, 1989, 1993; Kojima et al. 1993), Winnepenninckx et al. suggested that pogonophorans and vestimentiferans are related to echiurans. For ease of discussion, we will assume that the vestimentiferans are within the pogonophoran clade, unless otherwise stated.

Evolutionary relationships within the Vestimentifera also are poorly resolved. Twelve species are assigned to eight genera and seven families. The lack of shared-derived characters between taxa has led to a preponderance of monogeneric families (six) and monospecific genera (six). Furthermore, some taxa exhibit extraordinary morphological plasticity that has misled several investigators to propose new species that did not withstand the scrutiny of subsequent molecular analyses. For example, two morphotypes of the genus *Ridgeia* were described as distinct species (Jones 1985), and six additional morphospecies were suspected from vents along the Juan de Fuca/Gorda/Explorer Ridge system (Tunnicliffe 1983, 1988; Tunnicliffe et al. 1985, 1986). Nevertheless, an allozyme survey detected no evidence for reproductive isolation or substantive geographical subdivision among these morphotypes (Black 1991). These morphotypes are now included in a single polytypic species, *R. piscesae* (Southward et al. 1995). Morphological plasticity also occurs in the polytypic vent tubeworm *Riftia pachyptila* (Black et al. 1994). Recognition of this plasticity does not preclude, however, the existence of cryptic species in other tubeworm taxa. For example, cryptic species are common among vent- and seep-endemic vesicomid clams (Vrijenhoek et al. 1994; Peek et al. 1997).

Questions also exist concerning the evolutionary age of vestimentiferan tubeworms, as these soft-bodied animals have left a limited and ambiguous fossil record. Some phylogenetically ancient relicts, notably stalked barnacles and slit limpets, flourish at vents, and thus have led some investigators to hypothesize that vent

habitats served as refugia for ancient lineages that otherwise failed to survive mass extinctions in the photic zone (McLean 1985; Newman 1985). A well-preserved Silurian vent community with fossilized tubes was thought to contain vestimentiferans (Little et al. 1997). Fossilized "tubeworm assemblages" were reported from fossilized vent sites as old as the mid to late Carboniferous (~325 million years ago), but these tubes may have been left by a variety of marine invertebrates, ranging from polychaetes to holothurians (Tunnicliffe 1991). Less controversial reports of fossilized vestimentiferan tubeworms are from the late Cretaceous Samail ophiolites in Oman (Haymon and Sinclair 1984; Haymon and Koski 1985).

Our goals in this study were: (1) to elucidate higher-level evolutionary relationships between vestimentiferan tubeworms and other protostome taxa; (2) to characterize genetic diversity and evolutionary relationships among vestimentiferan lineages; and (3) to assess the evolutionary age of vestimentiferans as a group. To accomplish these goals, we provide new DNA sequences from vestimentiferan specimens representing eight species and six genera. Representatives of four protostome phyla were used as outgroups for the higher level phylogenetic comparisons. All analyses were based on a 650 base pair (bp) region of the mitochondrial cytochrome oxidase subunit I gene (COI).

Materials and methods

Specimens examined in this study were collected during several oceanic cruises (1992, 1994 and 1995) by one or more of the authors, and the collections and voucher specimens are stored at Rutgers University. Several of the outgroup taxa were purchased as living specimens from biological supply companies (see Table 1). Specimens were either frozen at -80°C immediately upon collection and DNA was extracted subsequently at Rutgers University, or DNA was extracted from fresh tissue and the remaining carcasses are in -80°C storage. The two *Ridgeia* specimens from the Southern Explorer Ridge (the ridge system immediately to the north of the Juan de Fuca Ridge and separated by the Sovanco Transform fault) were collected in July of 1994. One specimen came from a collection that falls within the known range of morphological variation for *R. piscesae* (Jones), while the other came from an adjacent collection which did not. This second morphotype is tentatively referred to as *Ridgeia* sp. until the collection can be analyzed in greater detail. *Escarpia spicata* (Jones) was collected at a cold-water seep site and a hydrothermal vent site in the Guaymas Basin in May 1995. All vestimentiferan specimens were identified morphologically by MBB prior to DNA analyses. The identification of *Galathea alinum brachiosum* (Ivanov) was made by Dr. E. Southward at the time of collection, and later confirmed in her laboratory.

Molecular methods

Total genomic DNA was extracted using a modified hexadecyltrimethyl-ammonium bromide (CTAB) protocol (Doyle and Dickson 1987). A ≈ 710 base pair region of the mitochondrial cytochrome c oxidase subunit I gene was amplified via PCR using the primers LCO1490 and HCO2198 described in Folmer et al. (1994). The 50 μl PCR (polymerase chain-reactions) consisted of 35 to

70 ng of template DNA, 5 µl of the 10× buffer supplied by the manufacturer, 2.5 mM MgCl₂, 200 µM dNTPs, 0.5 µM of each primer, and 1.5 U *Taq* polymerase (Promega Inc., Wisconsin). After an initial denaturation at 95 °C for 2 min, reactions were amplified during 35 cycles as follows: 1 min at 95 °C, 1 min at 40 °C, and 1.5 min at 72 °C, followed by a final extension step of 7 min at 72 °C. For each set of PCR reactions, negative controls and sterile laboratory techniques were employed. PCR products were isolated with 1.5% NuSieve (FMC Bioproducts, New Jersey) agarose gels stained with ethidium bromide. The gel-isolated products were purified using either the Wizard-PCR kit (Promega, Wisconsin) or the Qiagen gel extraction kit (Chatsworth, California).

The sequences reported here were originally obtained with “manual” sequencing methods. However, due to the presence of some ambiguities, all sequences (except the *Riftia pachyptila* – 9°N, East Pacific Rise specimen) were verified by automated sequencing. In both cases, the LCO1490 and HCO2198 oligonucleotides served as sequencing primers.

For the manually obtained sequences, 55 ng to 80 ng purified PCR product was sequenced with the AmpliTaq cycle sequencing kit (Perkin-Elmer Cetus, California). The cycle-sequencing parameters were as follows: 20 cycles of 95 °C for 1 min, 65 °C for 1 min, 72 °C for 1.5 min. The primers were end-labeled with γ³³P to allow visualization of fragments that were separated by a 5 to 6% denaturing polyacrylamide gel. For each primer, short- and long-sequence reads were combined and light- and heavy-strand reads compared using the sequence editor Gene Runner (Version 3.0, Hastings Software, Hastings, New York).

The automated sequence reactions followed the manufacturer’s (FS Dye Termination Mix, Applied Biosystems Inc., California) recommendations using ≈60 ng of purified PCR product as template. The reaction profile was 25 repetitions of denaturation at 96 °C for 30 s annealing at 50 °C for 15 s, and extension at 64 °C for 4 min. Fragments were visualized by use of dye-labeled terminator nucleotides. The samples were electrophoretically separated on a Perkin Elmer ABI 373A DNA sequencer. Data were analyzed using the AutoAssembler and Sequence Navigator programs (Applied Biosystems Inc., California). Both strands of the PCR product were completely sequenced. A comparison of the manually obtained sequences to the automated sequences allowed for the resolution of the ambiguous sites.

Analyses

Table 1 lists the operational taxonomic units (OTUs) examined, their collection locality, the identifying label used throughout this paper, and their GenBank accession number. Final sequences were entered into the Eyeball Sequence Editor editor (ESEE; Cabot and Beckenbach 1989), and aligned using the amino acid sequences of the following reference taxa: *Drosophila yakuba* (GenBank Accession X03240; Clary and Wolstenholme 1985), *Daphnia pulex* (GenBank Accession Z15015; Van Raay and Crease 1994), *Anopheles gambiae* (GenBank Accession L20934; Beard et al. 1993) and *Artemia franciscana* (GenBank Accession X69067; Perez et al. 1994). The alignment consisted of 650 nucleotide positions, corre-

Table 1 Taxa used in analysis of a 710 base pair fragment of mitochondrial cytochrome oxidase subunit I gene (*OTU* operational taxonomic unit)

Organism	Source	OTU label	GenBank accession
Vestimentifera			
Axonobranchia			
<i>Riftia pachyptila</i>	9°N Eastern Pacific Rise (09°31'N, 104°18'W; 2564 m)	<i>Rpa</i> 9N	U74074
<i>Riftia pachyptila</i>	18°S Eastern Pacific Rise (18°36'S, 113°24'W; 2673 m)	<i>Rpa</i> 18S	U74053
Basibranchia			
<i>Escarpia laminata</i>	Florida Escarpment (26°02'N, 84°54'W; 3310 m)	<i>Ela</i>	U74063
<i>Escarpia spicata</i>	Guaymas Basin seep (27°35'N, 111°28'W; 1653 m)	<i>Esp</i> seep	U74065
<i>Escarpia spicata</i>	Guaymas Basin vent (27°00'N, 111°25'W; 2020 m)	<i>Esp</i> vent	U74064
<i>Lamellibrachia barhami</i>	Middle Valley (48°27'N, 128°42'W; 2416 m)	<i>Lba</i> MV	U74055
<i>Lamellibrachia barhami</i>	Oregon Margin (44°41'N, 125°17'W; 2089 m)	<i>Lba</i> OM	U74054
<i>Lamellibrachia columna</i>	Lau Basin (22°32'S, 176°43'W; 1870 m)	<i>Lco</i>	U74061
<i>Oasisia alvinae</i>	21°N Eastern Pacific Rise (20°47'N, 109°09'W; 2577 m)	<i>Oal</i>	U74069
<i>Ridgeia piscesae</i>	Juan de Fuca Ridge (44°39'N, 130°22'W; 2227 m)	<i>Rpi</i> SJR	U74073
<i>Ridgeia piscesae</i>	Southern Explorer Ridge (49°46'N, 130°16'W; 1780 m)	<i>Rpi</i> SER	U74057
<i>Ridgeia</i> sp.	Southern Explorer Ridge (49°46'N, 130°16'W; 1780 m)	<i>Rsp</i> SER	U74056
<i>Tevnia jerichonana</i>	9°N Eastern Pacific Rise (09°50'N, 104°17'W; 2550 m)	<i>Tje</i>	U74075
Undescribed species	Nikko Seamount (23°04.6'N, 142°19.8'E; 433 m)	Nikko vest.	U74078
Pogonophora			
<i>Galatheolinum brachiosum</i>	Oregon Margin (45°01'N, 125°17'W; 2028 m)	<i>Gbr</i>	U74066
Undescribed species	Loihi Seamount (18°55'N, 155°15'W; 915 m)	Lohi pogo.	U74068
Annelida			
Hirudinea			
<i>Hirudo medicinalis</i>	Purchased from Carolina Biological Supply Co.	<i>Hme</i>	U74067
Oligochaeta			
<i>Tubifex tubifex</i>	Purchased from Carolina Biological Supply Co.	<i>Ttu</i>	U74076
Polychaeta			
<i>Paralvinella palmiformis</i>	Juan de Fuca Ridge (47°58.2'N, 129°05.2'W; 2180 m)	<i>Ppa</i>	U74070
Echiura			
<i>Urechis</i> sp.	Purchased from Pacific Bio Marine Supply Co.	<i>Urechis</i> sp.	U74077
Sipuncula			
Phascolosomida			
<i>Phascolosoma</i> sp.	Purchased from Pacific Bio Marine Supply Co.	<i>Phascolosoma</i> sp.	U74071

sponding to 216 amino acid residues which contained only one amino acid deletion in *D. pulex*. The inferred amino acid sequences were obtained using the *Drosophila* mitochondrial translation code.

Phylogenetic analyses were performed using maximum parsimony (MP), maximum-likelihood (ML), and neighbor-joining (NJ) for both nucleotide and amino acid residues. PAUP (phylogenetic analysis using parsimony; Version 3.1.1; Swofford 1993) was used to conduct the parsimony analyses. Branch-and-bound searches were used to find the best tree(s) for the nucleotide data set, while heuristic searches were used in all parsimony bootstrap-analyses and to find the best amino acid tree. Neighbor-joining analysis was completed using the phylogeny inference (PHYLIP) package (Version 3.5c; Felsenstein 1993) and employed a Kimura two-parameter correction model. The nucleotide likelihood analyses were conducted using fastDNAML (Olsen et al. 1994) with a Kimura two-parameter correction. Protein maximum likelihood (PROTML; Version 2.2; Adachi and Hagesawa 1992) was used to obtain the likelihood tree for the amino acid data based on a Jones-Taylor-Thornton (Jones et al. 1992) amino acid substitution matrix. Bootstrap analysis for both parsimony and neighbor-joining consisted of 500 iterations, whereas the likelihood analysis only employed 100 iterations (due to computation time limitations). Computation time and software limitations made a likelihood bootstrap of the amino acid data prohibitive.

Because of the problem of third-position transitional saturation of protein coding mitochondrial genes (Brown et al. 1982; Graybeal 1993; Adkins and Honeycutt 1994; Yoder et al. 1996), the MacClade (Version 3.06; Maddison and Maddison 1992) software package was used to recode third-position nucleotides to remove transitional state changes prior to all analyses of the nucleotide data.

Results

Our aim was to obtain as complete a representation as possible of the Vestimentifera. The following taxa were not available due to their rarity: the undescribed Vigo worm (Dando et al. 1992; Williams et al. 1993), *Alaysia spiralis* (Southward 1991), *Lamellibrachia luymsi* (Van der Land and Nørrevang 1975) and *L. victori* (Mañé-Garzón and Montero 1985). While pogonophoran species have been reported from all the world oceans, they are often found in remote regions and thus are not easily collected.

Due to the uncertainty of vestimentiferan/pogonophoran relationships (Kojima et al. 1993; Winnepeinckx et al. 1995), we examined a variety of protostome taxa to determine appropriate outgroup taxa. First, we reconstructed these higher level relationships by taking advantage of the conservative nature of amino acid substitutions in the COI gene. Subsequent analyses focusing on the ingroup used the original nucleotide data, with third positions recoded to remove transitions.

Of the 216 inferred amino acid (a.a.) positions, 118 were variable and 81 were informative (with respect to parsimony analysis) across the full range of protostomes examined. However, within the Vestimentifera, we found only one parsimony-informative a.a. substitution (Position 139) which suggests that *Riftia* and *Tevnia* are sister taxa. Although different outgroup topologies were obtained with maximum parsimony (MP), maximum likelihood (ML), and neighbor-joining (NJ) methods, all reconstructions placed annelids as the closest pogonophoran relatives.

Fig. 1a shows the maximum-likelihood reconstruction obtained from these a.a. sequences (log likelihood score = -2152.19). Based on previously published phylogenies (e.g. Eernisse et al. 1992; Halanych et al. 1995), we rooted the topology along the branch separating arthropods from the remaining protostome taxa. The results of bootstrap analyses are shown on the topology. Although we prefer ML as a method of phylogeny estimation (see Swofford et al. 1996), the NJ topology (Fig. 1b) is more consistent with the current understanding of protostome phylogeny. For example, the two crustaceans are monophyletic. Parsimony analysis (heuristic search option with tree-bisection-reconnection branch swapping) identified 148 "shortest" trees (369 steps; confidence interval, CI = 0.797, CI minus uninformative characters = 0.601). Most variation among these trees was due to the zero-length branches within the vestimentiferan clade, and the strict consensus of these trees was identical to the ML reconstruction except that the echiurid (*Urechis* sp.) and the scipunculan (*Phascolosoma* sp.) fell out as paraphyletic with the sipunculan more basal. All three methods were similar with respect to placement of the pogonophoran clade and the polychaete annelid *Paralvinella* sp. at the base of the pogonophoran clade.

Based on these amino acid reconstructions, the taxa employed in the analyses of nucleotide data were trimmed to include the Pogonophora and the three annelid taxa as outgroups (19 total taxa). The nucleotide data included 287 variable positions and 202 parsimony-informative sites when third-position transitions were excluded. These values fell to 164 variable and 111 informative sites upon exclusion of the annelid outgroups.

All three methods of phylogeny reconstruction (MP, ML, and NJ) recovered the same topology suggesting that the signal in the data is robust under a variety of assumptions. This topology (shown in Fig. 2) includes proportional branch lengths obtained by likelihood reconstruction (log likelihood score = -5861.38). The most parsimonious tree obtained with the PAUP branch and bound algorithm consisted of 585 steps, with a CI of 0.568 (CI minus uninformative characters = 0.484). The bootstrap values for all three reconstruction methods are shown in Fig. 2. In general, the tips of the tree were well supported, whereas deeper nodes within the Pogonophora had lower bootstrap support.

Distance values based on a Kimura two-parameter correction of the nucleotide data are shown in Table 2. For the purpose of comparison with published data, the reported distance values were based on the original nucleotide data (i.e. not recoded third positions). The number of absolute differences between OTUs is also shown.

Discussion

The present results with mitochondrial COI sequences indicate that vestimentiferan tubeworms are nested


Fig. 1 Trees based on cytochrome c oxidase subunit I, inferred amino acid sequences (216 residues). **a** Protein maximum-likelihood tree (PROTML Version 2.2) with branch lengths estimated from a Jones–Taylor–Thornton substitution matrix (1992) and a STAR decomposition search; parsimony bootstrap values (500 iterations, see “Results”) and a scale bar of branch length as percent probability of change along a given branch are shown; unfortunately, due to computation time and software limitations, a maximum-likelihood bootstrap analysis was not possible. **b** Neighbor-joining tree based on a PAM matrix; branch lengths represent corrected amino acid distances calculated with ProtDIST program in PHYLIP package (*scale bar* = percent change) (*Numbers* proportion of occurrence in 500 neighbor-joining replicates, bootstrap values of < 60% are not shown for either **a** or **b**; OTU labels as in Table 1)



within the Pogonophora. Recent analysis of nuclear small-subunit (18S) rDNA sequences also identified a close relationship between perviate pogonophorans and vestimentiferans, and supported inclusion of the two taxa in a single clade (Winnepenninckx et al. 1995). However, basing their findings on the nuclear 18S rDNA, Winnepenninckx et al. suggested that pogonophorans may be more closely related to echiurans than annelids. The present phylogenetic reconstructions

based on COI indicate that pogonophorans share an evolutionary history with annelids, and place echiurans outside this group (Fig. 1). In all our reconstructions, one or more of the annelid taxa (typically the polychaete *Paralvinella palmiformis*) occurred immediately basal to the Pogonophora. This placement is consistent with other phylogenetic analyses (Bartolomaeus 1995; Rouse and Fauchald 1995; Malakhov et al. 1996; McHugh 1997).

Fig. 2 Tree based on cytochrome c oxidase subunit I, DNA sequences (650 base pairs – recoded to remove third codon position transitions; OTU labels as in Table 1). Topology obtained with parsimony (PAUP, branch and bound search, TBR branch-swapping), maximum-likelihood (PHYLIP, Kimura two-parameter model), and neighbor-joining (PHYLIP, Kimura two-parameter model) reconstruction algorithms [Numbers next to given node percentage bootstrap support: plain numbers likelihood values out of 100 iterations; underlined numbers parsimony values out of 500 iterations; italicised numbers neighbor-joining values with 500 iterations (bootstrap values < 60% are not shown); values between OTUs for *Ridgeia piscesae* and *Escarpia spicata* indicate bootstrap support for respective taxa; however, branch lengths in these situations are extremely short and, to some degree, values may result from strictly bifurcating assumptions made by “neighbor” and “fast-DNAml boot” programs



Presently, the higher-order relationships between Annelida, Pogonophora and other protostome invertebrates remain ambiguous. The paraphyletic relationship between the pogonophoran taxa included in this study is not without precedent. Based on morphological criteria, Rouse and Fauchald (1995) discussed the possible paraphyly of the frenulate pogonophorans and the Polychaeta. Monophyly of the Annelida seems increasingly unsupported, as is evidenced by the present results. Nevertheless, a problem with the present analysis and all other molecular studies to date is a strong representational bias regarding the taxa included for phylogenetic analyses. The Annelida comprise > 14 000 species and the Pogonophora > 140 species. The representation of these groups in molecular analyses has been extremely limited, leading perhaps to biases that may have profound effects on phylogenetic reconstructions of these diverse groups (see Swofford and Olsen 1990).

Unfortunately, the fossil record for soft-bodied worms is notoriously poor and incomplete (see Bengtson 1994 and other studies cited therein), and thus it is difficult to make reasonable guesses concerning divergence times for these taxa based on morphological grounds. However the molecular data suggest that vestimentiferans appear to constitute a recent group compared to the outgroup taxa (Fig. 1). Furthermore, the short internal and terminal branches within the vestimentiferan clade suggest that the present distribution of taxa can be accounted for by a recent rapid radiation. COI is one of the most conservative genes in the mitochondrial genome with respect to amino acid substitution. We found < 1% sequence divergence among the extant vestimentiferans. According to the estimates of COI a.a. sequence divergence presented by Wray et al. (1996), one interpretation of our data is that the Vestimentifera as a taxon may be < 100 million years old, indicating a more recent divergence than is traditionally recognized.

Table 2 Pairwise distance comparisons for nucleotide data (including third-position transitions) (Above diagonal genetic distances corrected with the Kimura two-parameter model; below diagonal absolute pairwise differences) OTU labels and full specific names in Table 1

	Rpa 9N	Rpa 18S	Ela	Exp seep	Exp vent	Lba MV	Lba OM	Lco	Oal	Rpi SJR	Rpi SER	Rsp SER	Tje	Nikko vest.	Gbr	Loihi pogo	Hme	Ttu	Ppa
Rpa 9N																			
Rpa 18S	0																		
Ela	107	109																	
Exp seep	109	111	2																
Exp vent	108	110	1	1															
Lba MV	100	102	91	91	90														
Lba OM	102	104	92	92	91	2													
Lco	103	106	90	90	89	31	31												
Oal	96	97	89	89	88	93	95	91											
Rpi SJR	97	99	108	108	107	93	95	93	80										
Rpi SER	96	98	106	106	105	92	94	92	77	3									
Rsp SER	96	99	107	107	106	92	94	92	77	3	0								
Tje	86	87	84	86	85	93	96	93	92	88	88	88							
Nikko vest.	103	105	87	87	86	30	30	33	97	96	95	95	93						
Gbr	133	138	130	129	129	125	127	126	134	131	133	133	132	118					
Loihi pogo.	123	124	113	111	112	111	114	113	118	118	118	118	113	112	141				
Hme	207	211	214	214	213	209	212	215	208	213	213	213	213	203	176	213			
Ttu	185	189	184	182	183	181	182	179	180	188	190	191	180	175	186	198	209		
Ppa	179	183	182	184	183	170	171	174	184	183	184	184	170	165	173	184	198	180	

Alternatively, one can argue that the rate of evolutionary change of the COI gene within the Vestimentifera has slowed substantially. Because this estimate is based on a single gene sequence, the evolutionary interpretation must be viewed with caution. Currently, we are focusing on additional genetic markers to help resolve this issue of vestimentiferan age.

The present COI nucleotide-sequence data provides good resolution of relationships among vestimentiferan taxa (Fig. 2). The illustrated topology was fully supported by all three types of phylogenetic analyses (ML, NJ, and MP). Genera associated with eastern Pacific hydrothermal vents formed a monophyletic clade with two groups of sister taxa: *Oasisia* and *Ridgeia* and *Riftia* and *Tevnia*. The genera typically associated with cold-water sulfide/hydrocarbon seeps [*Lamellibrachia* (including the undescribed species from Nikko Seamount) and *Escarpia*] were paraphyletic and basal to the vent clade. Although the clustering of *Tevnia* with *Riftia* only received marginal bootstrap support, the monophyly of the *Oasisia*–*Ridgeia* clade is strongly supported by bootstrap values of 93, 84 and 99% for ML, MP and NJ analyses, respectively. The present relationships are different from the tree generated by Williams et al. (1993) based on 322 bp of the nuclear large-subunit (28S) rDNA. Their 28S rDNA data placed *Ridgeia* and *Tevnia* as sister taxa, with *Riftia* basal to the “vent” clade; however, they did not include *Oasisia* in their analysis. Furthermore, the assignment of higher taxonomic levels by Jones (1985) does not reflect genealogical relationships based on the COI sequences. His assignment of *Riftia pachyptila* to the monogeneric taxon *Axonobrachia*, based on arrangement of the branchial plume and its vascular system, is unsupported by the present data. In addition, our data do not support *Oasisia* and *Tevnia* as distinct from *Ridgeia*.

A major feature seen in the diversification of these species of vestimentiferan tubeworms was the association with specific kinds of deep-sea environments. Together, the genera *Riftia*, *Oasisia*, *Ridgeia*, and *Tevnia* form a clade that is endemic to hydrothermal vents of the eastern Pacific. In contrast, the genera *Escarpia* and *Lamellibrachia* are found in soft sediments associated with cold-water sulfide/hydrocarbon seeps. A recent molecular study (Feldman et al. 1997a) of bacterial endosymbionts associated with these vestimentiferans revealed a parallel, but not concurrent, subdivision of their bacteria (i.e. they did not coevolve). All species of vent-endemic tubeworms contain one endosymbiont species, while the vent-associated tubeworms contain another species which is markedly genetically different.

The two seep-associated genera are paraphyletic and basal to the vent-clade. However, some members of the “seep” genera appear to be opportunistic, also occurring in sedimented basins near hydrothermal vents. For example, *Escarpia spicata* is found near the Guaymas Basin hydrothermal vents in the Gulf of California, and *Lamellibrachia barhami* occurs near hydrothermal vents in Middle Valley on the Juan de Fuca Ridge (Black et al.

1997). Kojima et al. (1997) provide similar examples for *Escarpia* and *Lamellibrachia* in the vicinity of Japan. The best evidence for the opportunistic nature of seep-associated tubeworms was the recent discovery of *E. spicata* living in sulfide-rich sediments associated with a whale carcass (Feldman et al. 1997b). Perhaps the opportunistic nature of some seep-associated tubeworms may account for their invasion of hydrothermal areas. We suspect the present vent-endemic clade arose from such an opportunistic ancestor. This evolutionary pathway is also found in deep-sea mussels (Craddock et al. 1995). Seep-associated species are basal and paraphyletic, whereas hydrothermal vent-endemic species form a monophyletic and derived group. It will be interesting to see if this pattern of evolution is repeated in other groups such as vesicomyid clams and bresiliid shrimp.

Acknowledgements We thank the ship and submersible crews of the "Atlantis II/Alvin", "Nadir/Nautile", and the "Yokosuka/Shinkai 6500" for their hospitality and assistance. Our thanks to E. Southward who was fortuitously onboard during the collection of our sole *Galathealinum brachiosum* specimen, and to C. Moyer of the University of Hawaii for providing specimens. A. Trivedi and C. Di Meo assisted with PCR and sequencing. This is Contribution No. 97-09 of the Institute of Marine and Coastal Sciences, and New Jersey Agricultural Experiment Station Publication No. D-32104-5-97, supported by state funds and by NSF Grants OCE89-17311, OCE92-17026, and OCE93-02205.

References

- Adachi J, Hasegawa M (1992) MOLPHY: programs for molecular phylogenetics I – PROTML: maximum likelihood inference of protein phylogeny. Comput Sci Monogr (Tokyo) No. 27
- Adkins RM, Honeycutt RL (1994) Evolution of the primate cytochrome c oxidase subunit II gene. J molec Evolut 38: 215–231
- Bartolomaeus T (1995) Structure and formation of the uncini in *Pectinaria koreni*, *Pectinaria auricoma* (Terebellida) and *Spirorbis spirorbis* (Sabellida): implications for annelid phylogeny and the position of the Pogonophora. Zoomorphology 115: 161–177
- Beard CB, Hamm DM, Collins FH (1993) The mitochondrial genome of the mosquito *Anopheles gambiae*: DNA sequence, genome organization, and comparisons with mitochondrial sequences of other insects. Insect molec Biol 2: 103–124
- Bengtson S (1994) The advent of animal skeletons. In: Bengtson S (ed) Early life on Earth: Nobel Symposium No. 84. Columbia University Press, New York, pp 412–425
- Black MB (1991) Genetic (allozyme) variation in Vestimentifera (*Ridgeia* spp.) from hydrothermal vents of the Juan de Fuca Ridge (Northeast Pacific Ocean). University of Victoria, Victoria, British Columbia
- Black MB, Lutz RA, Vrijenhoek RC (1994) Gene flow among vestimentiferan tubeworm (*Riftia pachyptila*) populations from hydrothermal vents of the eastern Pacific. Mar Biol 120: 33–39
- Black MB, Trivedi A, Lutz RA, Vrijenhoek RC (1997) Population genetics and biogeography of vestimentiferan tube worms. Deep-Sea Res (in press)
- Brown WM, Prager Em, Wang A, Wilson AC (1982) Mitochondrial DNA sequences of primates: tempo and mode of evolution. J molec Evolut 18: 225–239
- Cabot EL, Beckenbach AT (1989) Simultaneous editing of multiple nucleic acid and protein sequences with ESEE. Cabios 5: 233–234
- Cavanaugh CM, Gardiner SL, Jones ML, Jannasch HW, Waterbury JB (1981) Prokaryotic cells in the hydrothermal vent tubeworm *Riftia pachyptila* Jones: possible chemoautotrophic symbionts. Science, NY 213: 340–342
- Clary DO, Wolstenholme DR (1985) The mitochondrial DNA molecule of *Drosophila yakuba*: nucleotide sequence, gene organization and genetic code. J molec Evolut 22: 252–271
- Craddock C, Hoeh WR, Gustafson RG, Lutz RA, Hashimoto J, Vrijenhoek RC (1995) Evolutionary relationships among deep-sea mytilids (Bivalvia: Mytilidae) from hydrothermal vents and cold-water methane/sulfide seeps. Mar Biol 121: 477–485
- Dando PR, Southward AJ, Southward EC, Dixon DR, Crawford A, Crawford M (1992) Shipwrecked tubeworms. Nature, Lond 356: p. 667
- Doyle JJ, Dickson E (1987) Preservation of plant samples for DNA restriction endonuclease analysis. Taxon 36: 715–722
- Eernisse DJ, Albert JS, Anderson FE (1992) Annelida and Arthropoda are not sister taxa: a phylogenetic analysis of spiralian metazoan morphology. Syst Biol 41: 305–330
- Felbeck H (1981) Chemoautotrophic potential of the hydrothermal vent tubeworm, *Riftia pachyptila* Jones (Vestimentifera). Science, NY 213: 336–338
- Feldman RA, Black MB, Cary CS, Lutz RA, Vrijenhoek RC (1997a) Molecular phylogenetics of bacterial endosymbionts and their vestimentiferan hosts. Molec Mar Biol Biotechnol 6: 268–277
- Feldman RA, Shank TM, Black MB, Baco A, Smith CR, Vrijenhoek RC (1997b) Vestimentiferan on a whale fall. Biol Bull mar biol Lab, Woods Hole 6: 268–277
- Felsenstein J (1993) PHYLIP (phylogeny inference package). Version 3.5c. Department of Genetics, University of Washington, Seattle, Washington
- Field KG, Olsen GJ, Lane DJ, Giovannoni SJ, Ghiselin MT, Raff EC, Pace NR, Raff RA (1988) Molecular phylogeny of the animal kingdom. Science, NY 239: 748–753
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molec mar Biol Biotechnol 3: 294–299
- Graybeal A (1993) The phylogenetic utility of cytochrome b: lessons from bufonid frogs. Mol Phylogenetics Evolut 2: 256–269
- Halanych KM, Bachelor J, Aquinaldo AM, Liva S, Hillis DM, Lake JA (1995) 18S rDNA evidence that the lophophorates are protostome animals. Science, NY 267: 1641–1643
- Haymon RM, Koski RA (1985) Evidence of an ancient hydrothermal vent community: fossil worm tubes in Cretaceous sulfide deposits of the Samail Ophiolite, Oman. Bull biol Soc Wash 6: 57–65
- Haymon RM, Sinclair C (1984) Fossil of hydrothermal vent worms from Cretaceous sulfide ores of the Samail Ophiolite, Oman. Science, NY 223: 1407–1409
- Ivanov AV (1988) Analysis of the embryonic development of Pogonophora in connection with the problems of phylogenetics. Zool Syst EvolForsch 26: 161–185
- Ivanov AV (1994) On the systematic position of Vestimentifera. Zool Jb (Abt Syst Geogr Biol Tiere) 121: 409–456
- Jones DT, Taylor WR, Thornton JM (1992) The rapid generation of mutation data matrices from protein sequences. Cabios 8: 25–282
- Jones ML (1981) *Riftia pachyptila*, new genus, new species, the vestimentiferan worm from the Galapagos Rift geothermal vents (Pogonophora). Proc biol Soc Wash 93: 1295–1313
- Jones ML (1985) On the Vestimentifera, new phylum: six new species, and other taxa, from hydrothermal vents and elsewhere. Bull biol Soc Wash 6: 117–158
- Jones ML (1987) On the status of the phylum-name, and other names, of the vestimentiferan tubeworms. Proc biol Soc Wash 100: 1049–1050
- Jones ML, Gardiner SL (1989) On the early development of the vestimentiferan tubeworm *Ridgeia* sp. and observations on the nervous system and trophosome of *Ridgeia* sp. and *Riftia pachyptila*. Biol Bull mar biol Lab, Woods Hole 177: 254–276
- Kojima S, Hashimoto T, Hasegawa M, Murata S, Ohta S, Seki H, Okada N (1993) Close phylogenetic relationship between

- Vestimentifera (tubeworms) and Annelida revealed by the amino acid sequence of elongation factor-1a. *J molec Evolut* 37: 66–70
- Kojima S, Segawa R, Hashimoto J, Ohta S (1997) Molecular phylogeny of vestimentiferans collected around Japan, revealed by the nucleotide sequences of mitochondrial DNA. *Mar Biol* 127: 507–513
- Little CTS, Herrington RK, Maslennikov VV, Morris NJ, Zaykov VV (1997) Silurian hydrothermal-vent community from the southern Urals, Russia. *Nature, Lond* 385: 146–148
- Maddison WP, Maddison DR (1992) *MacClade*. Sinauer, Sunderland, Massachusetts
- Malakhov VV, Popelyaev IS, Galkin SV (1996) Microscopic anatomy of *Ridgeia phaeophiale* (Jones, 1985) (Pogonophora, Vestimentifera) and the problem of the position of the Vestimentifera in the system of animal kingdom. I. General anatomy, obturacula, and tentacles. *Russ J mar Biol* 22: 63–74 [Transl of Biol Morya]
- Mañé-Garzón F, Montero R (1985) Sobre una nueva forma de verme tubícola *Lamellibrachia victori* n. sp. (Vestimentifera). Proposición de un nuevo phylum: Mesoneurophora. *Revta biol Urug* 8: 1–28
- McHugh D (1997) Molecular evidence that echiurans and pogonophorans are derived annelids. *Proc natn Acad Sci USA* 94: 8006–8009
- McLean JH (1985) Preliminary report on the limpets at hydrothermal vents. *Bull biol Soc Wash* 6: 159–166
- Newman WA (1985) The abyssal hydrothermal vent invertebrate fauna, a glimpse of antiquity? *Bull biol Soc Wash* 6: 231–242
- Nørrevang A (1970) On the embryology of *Siboglinum* and its implications for the systematic position of the Pogonophora. *Sarsia* 42: 7–16
- Nørrevang A (1974) Photoreceptors of the phaosome (hirudinean) type in a Pogonophore. *Zool Anz* 193: 297–304
- Olsen GJ, Matsuda H, Hagstrom R, Overbeek R (1994) fast-DNAml: a tool for construction of phylogenetic trees of DNA sequences using maximum likelihood. *Cabios* 10: 41–48
- Peek AS, Gustafson RG, Lutz RA, Vrijenhoek RC (1997) Evolutionary relationships of deep-sea hydrothermal vent and cold-water seep clams (Bivalvia: Vesicomidae): results from mitochondrial cytochrome oxidase subunit I. *Mar Biol* 130: 151–161
- Perez ML, Valverde JR, Batuecas B, Amat F, Marco R, Garesse R (1994) Speciation in the *Artemia* genus: mitochondrial DNA analysis of bisexual and parthenogenetic bring shrimps. *J molec Evolut* 38: 156–168
- Rouse GW, Fauchald K (1995) The articulation of annelids. *Zoologica Ser* 24: 269–301
- Southward EC (1988) Development of the gut and segmentation of newly settled stages of *Ridgeia* (Vestimentifera): implications for relationship between Vestimentifera and Pogonophora. *J mar biol Ass UK* 68: 465–487
- Southward EC (1991) Three new species of Pogonophora, including two vestimentiferans, from hydrothermal sites in the Lau Back-arc Basin (Southwest Pacific Ocean). *J nat Hist* 25: 859–881
- Southward EC (1993) Pogonophora. In: Harrison FW, Rice ME (eds) *Microscopic anatomy of invertebrates*. Vol. 12. Onychophora, Chilopoda, and lesser Protostomata. Wiley-Liss, New York, pp 327–369
- Southward EC, Tunnicliffe V, Black M (1995) Revision of the species of *Ridgeia* from northeast Pacific hydrothermal vents, with a redescription of *Ridgeia piscesae* Jones (Pogonophora: Obturata = Vestimentifera). *Can J Zool* 73: 282–295
- Suzuki T, Takagi T, Ohta S (1988) N-terminal amino acid sequence of the deep-sea tubeworm haemoglobin remarkably resembles that of annelid haemoglobin. *Biochem J* 255: 541–545
- Suzuki T, Takagi T, Ohta S (1993) N-terminal amino acid sequences of 440 · kDa hemoglobins of the deep-sea tubeworms, *Lamellibrachia* sp. 1, *Lamellibrachia* sp. 2 and slender Vestimentifera gen. sp. 1. Evolutionary relationship with annelid hemoglobins. *Zool Sci* 10: 141–147
- Suzuki T, Takagi T, Okuda K, Furukohri T, Ohta S (1989) The deep-sea tubeworm hemoglobin: subunit structure and phylogenetic relationship with annelid hemoglobin. *Zool Sci* 6: 915–926
- Swofford DL (1993) PAUP: Phylogenetic analysis using parsimony. Version 3.1.1. Computer program distributed by the Illinois Natural History Survey, Champaign, Illinois, USA
- Swofford DL, Olsen GJ (1990) Phylogeny Reconstruction. In: Hillis DM, Moritz C (eds) *Molecular systematics*. Sinauer, Massachusetts, pp 411–501
- Swofford DL, Olsen GJ, Waddell PJ, Hillis DM (1996) Phylogenetic inference. In: Hillis DM, Moritz C (eds) *Molecular systematics*. Sinauer, Massachusetts, pp 407–514
- Tunnicliffe V (1983) New hot vents in NE Pacific. *EOS Trans, Am geophys Un* 64: p. 945
- Tunnicliffe V (1988) Biogeography and evolution of hydrothermal-vent fauna in the eastern Pacific Ocean. *Proc R Soc (Ser B)* 233: 347–366
- Tunnicliffe V (1991) The biology of hydrothermal vents: ecology and evolution. *Oceanogr mar Biol A Rev* 29: 319–407
- Tunnicliffe V, Botros M, De Burgh ME, Dinet A, Johnson HP, Juniper SK, McDuff RE (1986) Hydrothermal vents of Explorer Ridge, northeast Pacific. *Deep-sea Res* 33: 401–412
- Tunnicliffe V, Juniper SK, De Burgh ME (1985) The hydrothermal vent community on Axial Seamount, Juan de Fuca Ridge. *Bull biol Soc Wash* 6: 453–464
- Van der Land J, Nørrevang A (1975) The systematic position of *Lamellibrachia* [sic] (Annelida, Vestimentifera). *Z zool Syst EvolForsch (Sonderheft)*: 86–101
- Van Raay TJ, Crease TJ (1994) Partial mitochondrial DNA sequence of the crustacean *Daphnia pulex*. *Curr Genet* 25: 66–72
- Vrijenhoek RC, Schutz SJ, Gustafson RG, Lutz RA (1994) Cryptic species of deep-sea clams (Mollusca, Bivalvia, Vesicomidae) in hydrothermal vent and cold-seep environments. *Deep-Sea Res* 41: 1171–1189
- Webb M (1964) The larvae of *Siboglinum fiordicum* and a reconsideration of the adult body regions (Pogonophora). *Sarsia* 15: 57–68
- Webb M (1969) *Lamellibrachia barhami*, gen. nov., sp. nov. (Pogonophora), from the northeast Pacific. *Bull mar Sci* 19: 18–47
- Williams NA, Dixon DR, Southward EC, Holland PWH (1993) Molecular evolution and diversification of the vestimentiferan tubeworms. *J mar biol Ass UK* 73: 437–452
- Winnepenninckx B, Bäckeljau T, Wachter RD (1995) Phylogeny of protostome worms derived from 18S rRNA sequences. *Molec Biol Evolut* 12: 641–649
- Wray GA, Levinton JS, Shapiro LH (1996) Molecular evidence for deep Precambrian divergences among Metazoan phyla. *Science, NY* 274: 568–573
- Yoder AD, Vilgalys R, Ruvolo M (1996) Molecular evolutionary dynamics of cytochrome b in strepsirrhine primates: the phylogenetic significance of third-position transversions. *Molec Biol Evolut* 13: 1339–1350